

9	<b>Experimental details:</b>	
<b>a. Treatment</b>	:	<ol style="list-style-type: none"> <li>1. <i>Beauveria bassiana</i> commercial product (CFU: 1.0 X10<sup>8</sup>) @ 10g/kg seed</li> <li>2. <i>Beauveria bassiana</i> commercial product (CFU: 1.0 X10<sup>8</sup>) @ 20g/kg seed</li> <li>3. <i>Metarhizium anisopliae</i> commercial product (CFU: 1.0 X10<sup>8</sup>) @ 10g/kg seed</li> <li>4. <i>Metarhizium anisopliae</i> commercial product (CFU: 1.0 X10<sup>8</sup>) @ 20g/kg seed</li> <li>5. <i>Beauveria bassiana</i> commercial product (CFU: 1.0 X10<sup>8</sup>) @ 10g/kg seed + Diatomaceous earth @ 5g/kg seed</li> <li>6. <i>Beauveria bassiana</i> commercial product (CFU: 1.0 X10<sup>8</sup>) @ 20g/kg seed + Diatomaceous earth @ 5g/kg seed</li> <li>7. <i>Metarhizium anisopliae</i> commercial product (CFU: 1.0 X10<sup>8</sup>) @ 10g/kg seed +Diatomaceous earth @ 5g/kg seed</li> <li>8. <i>Metarhizium anisopliae</i> commercial product (CFU: 1.0 X10<sup>8</sup>) @ 20g/kg seed +Diatomaceous earth @ 5g/kg seed</li> <li>9. Deltamethrin 2.8 EC @ 0.04 ml/kg seed</li> <li>10. Untreated control</li> </ol>